



Extension Agricultural Weather Report


June 10, 2019 - June 17, 2019.

Weekly Temperature & Rainfall Data from FAWN Stations by District

Northwest District









- 7-day rainfall (inches)
- 7-day Maximum Temperature (°F) •
- 7-day Minimum Temperature (°F)




Southwest District




Southeast District



NWS 6-10 DAY OUTLOOKS



NWS 6-10 day Rainfall Outlook



NWS 6-10 day Temperature Outlook



STATEWIDE CROP SUMMARY

(Data provided by the National Agricultural Statistics Service Crop Progress and Condition Report)

General

There were 5.8 days suitable for field work.

Precipitation estimates ranged from trace amounts in multiple locations to 10.3 inches in Lacoochee (Pasco County).

The average temperature ranged from 75.1°F in Lake City (Columbia County) to 85.1°F in Curry Hammock State Park (Monroe County).

Livestock and Pastures

Pasture conditions continued to improve with recent rainfall.

Standing water in DeSoto County caused producers to continue looking for parasites in cattle.

Cattle condition remained mostly good.

Fruits and Vegetables

Watermelon harvest continued in Levy and Lafayette counties.

Abundant rainfall caused some concern about fruit quality.

Fungal and bacterial diseases were observed on bean plants in Bradford County.

A wide variety of fruits and vegetables continued to be planted and brought to market.

Field Crops

Some growers in Santa Rosa County replanted cotton and peanuts due to dry weather, as well as hog and deer damage.

Soybean planting was reported in Walton County. Rainfall was beneficial to sugarcane and rice in the southern peninsula.

Citrus

The citrus growing region cooled slightly, with daily highs in the mid 80s to low 90s.

La Belle (Hendry County) received the most rainfall, with 6.25 inches.

According to the June 13, 2019 U.S. Drought Monitor, the entire citrus growing region remained drought free.

Weekly Valencia orange harvest continued to wind down.

Most processing plants have closed for the season, and remaining fruit will be fresh squeezed.

Grove activities included spraying and young tree care.

Caretakers pushed old trees not considered viable for next season.

Next season's fruit continued developing, with golf ball sized Early oranges, ping pong ball sized Valencias, and baseball sized grapefruit.

Irrigation slowed in areas where rainfall was plentiful, but some regular irrigation was still reported in the eastern part of the region.

STATEWIDE SOIL MOISTURE SUMMARY

(Data provided by the National Agricultural Statistics Service Crop Progress and Condition Report)

TOPSOIL	THIS WEEK (%)	PREVIOUS WEEK (%)
Very Short	3	7
Short	18	27
Adequate	66	59
Surplus	13	7

