


Extension Agricultural Weather Report


June 03, 2019 - June 10, 2019.

Weekly Temperature & Rainfall Data from FAWN Stations by District


Northwest District



Central District




Northeast District




● 7-day rainfall (inches) ● 7-day Maximum Temperature (°F) ● 7-day Minimum Temperature (°F)




Southwest District




Southeast District



NWS 6-10 DAY OUTLOOKS



NWS 6-10 day Rainfall Outlook



NWS 6-10 day Temperature Outlook



STATEWIDE CROP SUMMARY

(Data provided by the National Agricultural Statistics Service Crop Progress and Condition Report)

General

There were 5.6 days suitable for field work. Precipitation estimates ranged from trace amounts in multiple locations to 8.39 inches in Pensacola (Escambia County). The average temperature ranged from 79.2°F in Jasper (Hamilton County) to 86.1°F in Curry Hammock State Park (Monroe County).

Livestock and Pastures

Pasture conditions improved slightly with recent rainfall. There was standing water reported in some areas, which caused producers to look for parasites in cattle. Cattle condition remained mostly good.

Fruits and Vegetables

Watermelon harvest was reported in Levy and Lafayette counties. A wide variety of fruits and vegetables continued to be planted and brought to market.

Field Crops

Much needed rainfall improved crop condition throughout the state, but caused hay cutting to slow or halt in Okeechobee and Jackson counties.

Citrus

The citrus growing region remained warm, with daily highs in the low to mid 90s in most locations, and upper 90s in a few areas. Significant rainfall occurred in a few isolated areas within the citrus growing region. Some locations received 3-4 inches, but most areas received 0.25 - 2.5 inches. According to the June 6, 2019 U.S. Drought Monitor, several northern counties experienced moderate drought conditions, but the overall citrus growing region remained drought free. Weekly Valencia orange harvest slowed as the season began to wind down, with less than 2 million boxes harvested for the week.

Most processing plants have closed for the season, with the remainder scheduled to finish processing this week or next.

Grove activities included mowing, spraying nutritionals and insecticides, fertilizing, herbicide application, and general grove maintenance. Next season's fruit was growing normally, with golf ball sized oranges and larger grapefruit. Irrigation was run several times per week in most areas.

STATEWIDE SOIL MOISTURE SUMMARY

(Data provided by the National Agricultural Statistics Service Crop Progress and Condition Report)

TOPSOIL	THIS WEEK (%)	PREVIOUS WEEK (%)
Very Short	7	26
Short	27	31
Adequate	59	38
Surplus	7	5

